

Underlined parts: Revisions made in October 2024

[Keep this information leaflet with this medication and be sure to read it before taking the medication.]

EVE A TABLETS EX

Class ② OTC drug

Antipyretic-Analgesic Medication Ibuprofen Combination Product for Menstrual Pain and Headaches

- Each dose (a total of two tablets) of **EVE A TABLETS EX** contains 200 mg of ibuprofen, an analgesic agent, to achieve an excellent analgesic effect. This product also contains allylisopropylacetylurea and anhydrous caffeine, which enhance the analgesic effect.

Precautions for Use



Contraindications

If you do not follow the precautions exactly as listed below, your current symptoms may worsen and you may be placed at increased risk of side effects or the risk of having some accidents.

- The following persons must not take this medication:**
 - Persons who have had an allergic reaction to any of the ingredients of EVE A TABLETS EX.
 - Persons who have experienced asthma after taking this medication or after taking other antipyretic-analgesics or common cold medications.
 - Children under 15 years of age.
 - Pregnant women whose due date is within 12 weeks ahead.
- Do not take any of the following medications during the period of taking EVE A TABLETS EX:**

Other antipyretic-analgesics, common cold medications, sedatives, and motion sickness medications
- Do not operate vehicles or machinery after taking this medication.**

(Sleepiness or such may occur.)
- This medication must not be taken soon before or soon after alcohol intake.**
- Do not use this medication for a long and continuing period.**



Consultation

- The following persons should consult a physician, dentist, pharmacist, or registered salesperson before taking this medication:**
 - Persons who are receiving any treatment from a physician or dentist.
 - Women who are or may be pregnant.
 - Breast-feeding women.
 - Elderly persons.
 - Persons who have had any allergic reactions to drugs or other substances.
 - Persons who have received a diagnosis of any of the following diseases: Heart diseases, kidney diseases, liver diseases, systemic lupus erythematosus, mixed connective tissue disease
 - Persons who have had any of the following diseases: Gastroduodenal ulcers, ulcerative colitis, Crohn's disease
- If any of the symptoms listed below develop after taking EVE A EX, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, dentist, pharmacist, or registered salesperson, presenting this leaflet.**

Affected area	Symptoms
Skin	Rash/redness, itching, bruise
Gastrointestinal	Nausea/vomiting, inappetence, stomach discomfort, stomachache, stomatitis, heartburn, heavy feeling in the stomach, gastrointestinal hemorrhage, abdominal pain, diarrhea, bloody stools
Neuropsychiatric	Dizziness
Cardiovascular	Palpitations
Respiratory	Shortness of breath
Other	Blurred vision, tinnitus, swelling, nasal bleeding, gingival bleeding, difficulty in stopping bleeding, bleeding, back pain, excessively decreased body temperature, general malaise

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of symptoms	Symptoms
Shock (anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness or such occurring immediately after taking this medication.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis	Symptoms, such as high fever, ocular hyperemia, eye discharge, sore lips, sore throat, rashes and redness over an extensive area of the skin, persisting or suddenly worsening.

®: Registered trademark

(Continued on the reverse side)

(Continued from the front side)

Name of symptoms	Symptoms
Hepatic impairment	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general malaise, inappetence or such occurring.
Renal disorders	Fever, rashes, decreased urine output, generalized swelling, general malaise, arthralgia (pain in joints), diarrhea or such occurring.
Aseptic meningitis	Symptoms such as severe headache with neck stiffness, fever, and nausea/vomiting occurring. (Frequent occurrences of such symptoms have been reported particularly among patients who are undergoing treatment for systemic lupus erythematosus or mixed connective tissue disease.)
Myocardial infarction	Pain with chest tightness, respiratory distress, and cold sweat.
Cerebrovascular disorders	The following symptoms suddenly occurring: Decreased or loss of consciousness, difficulty in moving hand and foot unilaterally, headache, vomiting, dizziness, difficulty speaking, and speech impairment.
Asthma	Breathing sounds, such as wheezing and whistling, respiratory distress and such occurring.
Aplastic anemia	The following symptoms occurring: Bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Sudden high fever, chills, sore throat or such occurring.

- Any of the conditions mentioned below may occur after taking EVE A TABLETS EX. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.
Constipation, sleepiness
- If there is no alleviation of the symptoms for which this medication is being taken, after 3 or 4 doses, stop taking this medication and consult a physician, dentist, pharmacist, or registered salesperson, presenting this leaflet.

Indications

- Relief of the following pains: Painful menses (menstrual pain), headache/shoulder stiffness pain, sore throat, joint pain, muscle pain, nerve pain, lower back pain, pain after tooth extraction, contusion pain, ear pain, fracture pain, sprain pain, and traumatic pain
- Relief of chills and fever


Dosage and Administration

Take a dose of EVE A TABLETS EX, according to the table below, with cold or lukewarm water, up to 2 times daily. Each dose should not be taken on an empty stomach whenever possible. There must be an interval of at least 6 hours between doses.

Age	Adults (aged 15 years or older)	Children under 15 years of age
Dose	2 tablets	× This medication must not be taken.

<Precautions concerning Dosage and Administration>

- (1) EVE A TABLETS EX must be used strictly in accordance with the specified dosage and administration.
- (2) How to remove the tablet: As shown in the illustration on the right, firmly press the raised portion of the press-through package (PTP) with a fingertip to push the tablet through the aluminum foil on the reverse side. (If the PTP sheet is mistakenly swallowed, its sharp edges may perforate the esophageal mucosa or lead to incidental events.)



Ingredients

in a total of 2 tablets

Ibuprofen	200 mg
Allylisopropylacetylurea	60 mg
Anhydrous caffeine	80 mg

Excipients: Anhydrous silicic acid, cellulose, croscarmellose sodium, polyvinyl alcohol-acrylic acid-methyl methacrylate copolymer, talc, magnesium stearate, hydroxypropylcellulose, hypromellose, macrogol, titanium oxide

Precautions for Storage and Handling

- (1) Store in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Do not take this medication after the expiration date.

Contact Information

Inquiries should be made to the drugstore where you purchased the product or to the customer service office.

Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193

Opening hours: 9:00 to 17:30 (closed on Saturdays, Sundays, and Japanese public holidays)

Manufactured and
Distributed by:



SSP Co., Ltd.

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1488

<https://www.ssp.co.jp/>